



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Abnormal enthalpy relaxation of GeO₂ in the merging region: does Johari-Goldstein relaxation exist in strong glasses?

Hu, Lina; Yue, Yuanzheng

Published in:
ICG 2007, XXI International Congress on Glass Abstract Book

Publication date:
2007

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Hu, L., & Yue, Y. (2007). Abnormal enthalpy relaxation of GeO₂ in the merging region: does Johari-Goldstein relaxation exist in strong glasses? In *ICG 2007, XXI International Congress on Glass Abstract Book* (pp. 205-205)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Abnormal enthalpy relaxation of germanium dioxides in the merging region: does Johari-Goldstein β relaxation exist in strong glasses?

Lina Hu, Yuanzheng Yue

Section of Chemistry, Aalborg University, Denmark

The merging of the primary α -relaxation and secondary relaxations (especially Johari-Goldstein β -relaxation), which occurs at a temperature somewhat above the standard glass transition temperature (T_g), has attracted much attention in recent years. For strong glasses with typical network structures, however, the secondary relaxation process, as well as the α - β merging region, is not well studied and understood due to the lack of a peak or a shoulder in the loss spectrum on the high frequency side of the primary α -relaxation. The present work focuses on the relaxation of the hyperquenched GeO_2 glasses. Upon fibre drawing, the GeO_2 glass is hyperquenched, and, hence, the liquid structure is frozen-in at a temperature well above T_g , but rather close to the liquidus temperature. Afterwards, the hyperquenched glass is subjected to annealing and subsequently to calorimetric scanning, by which information on enthalpy relaxation is obtained. The calorimetric results show that the enthalpy of the GeO_2 glass – one of the strongest glass systems, relaxes in quite a different manner than fragile systems. In the α - β merging region, all the primary α relaxation proceeds via the secondary relaxation. By using Williams's ansatz, we carried out further analysis on the dynamic properties of the secondary relaxation, in order to discover whether this secondary relaxation is real Johari-Goldstein β type or not. This study attempts to clarify whether the Johari-Goldstein β -relaxation is the intrinsic feature of the supercooled liquids and glasses. Such clarification is important for understanding glass transition phenomenon.